


Scale invariant fluid dynamics for the dilute Fermi gas at unitarity

Thomas Schaefer


North Carolina State University

Fluids: Gases, Liquids, Plasmas, . . .

Hydrodynamics: Long-wavelength, low-frequency dynamics of conserved or spontaneously broken symmetry variables.




$$\tau \sim \tau_{\text{micro}}$$



$$\tau \sim \lambda$$

Historically: Water
 $(\rho, \epsilon, \vec{\pi})$




Simple non-relativistic fluid

Simple fluid: Conservation laws for mass, energy, momentum

$$\frac{\partial \rho}{\partial t} + \vec{\nabla}(\rho \vec{v}) = 0$$

$$\frac{\partial \epsilon}{\partial t} + \vec{\nabla} \vec{j}^\epsilon = 0$$

$$\frac{\partial}{\partial t}(\rho v_i) + \frac{\partial}{\partial x_j} \Pi_{ij} = 0$$



Constitutive relations: Energy momentum tensor

$$\Pi_{ij} = P\delta_{ij} + \rho v_i v_j + \eta \left(\partial_i v_j + \partial_j v_i - \frac{2}{3} \delta_{ij} \partial_k v_k \right) + O(\partial^2)$$

reactive

dissipative

2nd order

$$\text{Expansion } \Pi_{ij}^0 \gg \delta \Pi_{ij}^1 \gg \delta \Pi_{ij}^2$$

Regime of applicability

Expansion parameter $Re^{-1} = \frac{\eta(\partial v)}{\rho v^2} = \frac{\eta}{\rho L v} \ll 1$

$$\frac{1}{Re} = \frac{\eta}{\hbar n} \times \frac{\hbar}{mvL}$$

fluid flow
property property


Consider $mvL \sim \hbar$: Hydrodynamics requires $\eta/(\hbar n) < 1$

Shear viscosity in kinetic theory

Kinetic theory: conserved quantities carried by quasi-particles

$$\frac{\partial f_p}{\partial t} + \vec{v} \cdot \vec{\nabla}_x f_p + \vec{F} \cdot \vec{\nabla}_p f_p = C[f_p]$$

$$\eta \sim \frac{1}{3} n \bar{p} l_{mfp}$$



Weakly interacting gas: $l_{mfp} \sim 1/(n\sigma)$ \Rightarrow $\eta \sim \frac{1}{3} \frac{\bar{p}}{\sigma}$

$$\eta(\sigma \rightarrow 0) \rightarrow \infty$$

Strongly interacting gas:

$$\frac{\eta}{n} \sim \bar{p} l_{mfp} \geq \hbar$$

but: kinetic theory not reliable!

Holographic duals at finite temperature



Thermal (conformal) field theory $\equiv AdS_5$ black hole

CFT temperature \Leftrightarrow

Hawking temperature of
black hole

CFT entropy \Leftrightarrow

Hawking-Bekenstein entropy
 \sim area of event horizon



$$s(\lambda \rightarrow \infty) = \frac{\pi^2}{2} N_c^2 T^3 = \frac{3}{4} s(\lambda = 0)$$

Gubser and Klebanov

Holographic duals: Transport properties

Thermal (conformal) field theory $\equiv AdS_5$ black hole

CFT entropy

\Leftrightarrow

Hawking-Bekenstein entropy

\sim area of event horizon

shear viscosity

\Leftrightarrow


Graviton absorption cross section

\sim area of event horizon

Strong coupling limit

$$\frac{\eta}{s} = \frac{\hbar}{4\pi k_B}$$


Son and Starinets (2001)



Strong coupling limit universal? Provides lower bound for all theories?

Answer appears to be no; e.g. theories with higher derivative gravity duals.

Effective theories for fluids (Unitary Fermi Gas, $T > T_F$)



$$\mathcal{L} = \psi^\dagger \left(i\partial_0 + \frac{\nabla^2}{2M} \right) \psi - \frac{C_0}{2} (\psi^\dagger \psi)^2$$



$$\frac{\partial f_p}{\partial t} + \vec{v} \cdot \vec{\nabla}_x f_p = C[f_p] \quad \omega < T$$



$$\frac{\partial}{\partial t}(\rho v_i) + \frac{\partial}{\partial x_j} \Pi_{ij} = 0 \quad \omega < T \left(\frac{T_F}{T} \right)^{3/2}$$

Effective theories (Strong coupling)




$$\mathcal{L} = \bar{\lambda}(i\sigma \cdot D)\lambda - \frac{1}{4}G_{\mu\nu}^a G_{\mu\nu}^a + \dots \Leftrightarrow S = \frac{1}{2\kappa_5^2} \int d^5x \sqrt{-g} \mathcal{R} + \dots$$

$$SO(d+2, 2) \rightarrow Schr(d) \qquad AdS_{d+3} \rightarrow \mathcal{X}_{d+3}$$




$$\frac{\partial}{\partial t}(\rho v_i) + \frac{\partial}{\partial x_j} \Pi_{ij} = 0 \quad (\omega < T)$$

Kinetics vs No-Kinetics



AdS/CFT low viscosity goo
gravitational dual

$$\eta/s \simeq 1/(4\pi)$$



kinetic liquid
quasi-particles


$$\eta/s \gtrsim 1$$

Outline

- I. Conformal second order hydrodynamics
- II. Kinetic theory
- III. Transport coefficients from elliptic flow

I. Non-relativistic fermions in unitarity limit

Consider simple square well potential




$$a < 0$$

$$a = \infty, \epsilon_B = 0$$

$$a > 0, \epsilon_B > 0$$

Non-relativistic fermions in unitarity limit

Now take the range to zero, keeping $\epsilon_B \simeq 0$



Universal relations


$$\mathcal{T} = \frac{1}{ik + 1/a}$$

$$\epsilon_B = \frac{1}{2ma^2}$$

$$\psi_B \sim \frac{1}{\sqrt{ar}} \exp(-r/a)$$

Universal fluid dynamics

Many body system: Effective cross section $\sigma_{tr} \sim n^{-2/3}$ (or $\sigma_{tr} \sim \lambda^2$)



Systems remains hydrodynamic despite expansion

Scale and conformal symmetry

Gallilean boosts $\vec{x}' = \vec{x} + t$ $t' = t$

scale trafo $\vec{x}' = e^s \vec{x}$ $t' = e^{2s} t$

conformal trafo $\vec{x}' = \vec{x}/(1 + ct)$ $1/t' = 1/t + c$

Ideal fluid dynamics

$$\Pi_{ij}^0 = P\delta_{ij} + \rho v_i v_j, \quad P = \frac{2}{3}\mathcal{E}$$

First order viscous hydrodynamics

$$\delta^{(1)}\Pi_{ij} = -\eta\sigma_{ij}, \quad \sigma_{ij} = \left(\nabla_i v_j + \nabla_j v_i - \frac{2}{3}\delta_{ij}(\nabla \cdot v) \right), \quad \zeta = 0$$

Scale invariant superfluid hydrodynamics

Momentum density: $\pi_i = \rho_n v_{n,i} + \rho_s v_{s,i}$

Stress tensor $\Pi_{ij} + \delta\Pi_{ij}$ with

$$\begin{aligned}\delta\Pi_{ij} = & -\eta \left(\nabla_i v_{n,j} + \nabla_j v_{n,i} - \frac{2}{3} \delta_{ij} \nabla_i v_{n,i} \right) \\ & - \delta_{ij} \left(\zeta_1 \nabla_i (\rho_s (v_{s,i} - v_{n,i})) + \zeta_2 (\nabla_i v_{n,i}) \right)\end{aligned}$$

Equation of motions for v_s : $\dot{v}_s + \frac{1}{2} \nabla (v_s^2) = -\nabla(\mu + H)$ with

$$H = -\zeta_3 \nabla_i (\rho_s (v_{s,i} - v_{n,i})) - \zeta_4 \nabla_i v_{n,i}$$

Conformal symmetry: $\zeta_1 = \zeta_2 = \zeta_4 = 0$


Why second order fluid dynamics?

Consider ideal expansion after release from a harmonic trap

$$\rho(x_i, t) = \rho_0(b_i(t)x_i), \quad v_i(x_j, t) = \alpha_i(t)x_i, \quad \alpha_i(t) = \dot{b}_i(t)/b_i(t)$$

Compare ideal and dissipative stresses

$$\Pi_{ii}^0 = P + \rho\alpha_i x_i^2, \quad \delta^{(1)}\Pi_{ii} = \eta(\alpha_i - \frac{4}{3}\sum_j \alpha_j)$$



Ideal stresses propagate with speed $\sim c_s$, dissipative stresses propagate with infinite speed. Hydro always breaks down in the dilute corona.

Second order conformal hydrodynamics

Relaxation of shear stress is a second order hydro term. Complete list

$$\begin{aligned}\delta^{(2)}\Pi^{ij} = & \eta\tau_\pi \left[\langle D\sigma^{ij} \rangle + \frac{2}{3}\sigma^{ij}(\nabla \cdot v) \right] \\ & + \lambda_1 \sigma^{\langle i}_k \sigma^{j\rangle k} + \lambda_2 \sigma^{\langle i}_k \Omega^{j\rangle k} + \lambda_3 \Omega^{\langle i}_k \Omega^{j\rangle k} \\ & + \gamma_1 \nabla^{\langle i} T \nabla^{j\rangle} T + \gamma_2 \nabla^{\langle i} P \nabla^{j\rangle} P + \gamma_3 \nabla^{\langle i} \nabla^{j\rangle} T + \dots\end{aligned}$$

$$A^{\langle ij \rangle} = \frac{1}{2} \left(A_{ij} + A_{ji} - \frac{2}{3} g_{ij} A^k{}_k \right) \quad \Omega^{ij} = (\nabla_i v_j - \nabla_j v_i)$$

New transport coefficients $\tau_\pi, \lambda_i, \gamma_i$

Can be written as a relaxation equation for $\pi^{ij} \equiv \delta\Pi^{ij}$

$$\pi^{ij} = -\eta\sigma^{ij} - \tau_\pi \left[\langle D\pi^{ij} \rangle + \frac{5}{3}(\nabla \cdot v)\pi^{ij} \right] + \dots$$

II. Linear response and kinetic theory

Consider background metric $g_{ij}(t, \mathbf{x}) = \delta_{ij} + h_{ij}(t, \mathbf{x})$. Linear response


$$\delta\Pi^{ij} = \frac{\delta\Pi_{ij}^{eq}}{\delta h_{ij}} h^{ij} - \frac{1}{2} G_R^{ijkl} h_{kl}$$

Kubo relation: $\eta(\omega) = \frac{1}{\omega} \text{Im}G_R^{xyxy}(\omega, 0)$

Kinetic theory: Boltzmann equation

$$\left(\frac{\partial}{\partial t} + \frac{p^i}{m} \frac{\partial}{\partial x^i} - \left(g^{il} \dot{g}_{lj} p^j + \Gamma_{jk}^i \frac{p^j p^k}{m} \right) \frac{\partial}{\partial p^i} \right) f(t, \mathbf{x}, \mathbf{p}) = C[f]$$

$$C[f] =$$




Kinetic theory

linearize $f = f_0 + \delta f$, solve for δf , $\hookrightarrow \delta \Pi_{ij}$, $\hookrightarrow G_R$, $\hookrightarrow \eta(\omega)$


$$\eta(\omega) = \frac{\eta}{1 + \omega^2 \tau_R^2}$$

$$\eta = \frac{15}{32\sqrt{\pi}}(mT)^{3/2} \quad \tau_R = \frac{\eta}{nT}$$

shear channel



sound channel



Shear viscosity: Sum rules



Randeria & Taylor proved the sum rules (corrected by Enss & Zwerger)

$$\frac{1}{\pi} \int dw \left[\eta(\omega) - \frac{C}{15\pi\sqrt{m\omega}} \right] = \frac{\mathcal{E}}{3} - \frac{C}{10\pi ma}$$

$$\frac{1}{\pi} \int dw \zeta(\omega) = \frac{1}{72\pi ma^2} \left(\frac{\partial C}{\partial a^{-1}} \right)$$

where C is Tan's contact, $\rho(k) \sim C/k^4$.

Sum rules constrain spectral function and euclidean correlator





Second order hydrodynamics from kinetic theory

Boltzmann equation (BGK approximation)


$$\begin{aligned}\delta^{(2)}\Pi^{ij} = & \frac{\eta^2}{P} \left[\langle D\sigma^{ij} \rangle + \frac{2}{3} \sigma^{ij} (\nabla \cdot v) \right] \\ & + \frac{\eta^2}{P} \left[\sigma^{\langle i}_k \sigma^{j\rangle k} + \sigma^{\langle i}_k \Omega^{j\rangle k} \right] + O(\kappa\eta\nabla^i\nabla^j T)\end{aligned}$$

relaxation time $\tau_R = \frac{\eta}{P} \simeq \frac{\eta}{nT}$


III. Almost ideal fluid dynamics (cold Fermi gas)




Hydrodynamic
expansion converts
coordinate space
anisotropy
to momentum space
anisotropy



Elliptic flow: High T limit




Quantum viscosity $\eta = \eta_0 \frac{(mT)^{3/2}}{\hbar^2}$



fit: $\eta_0 = 0.33 \pm 0.04$


theory: $\eta_0 = \frac{15}{32\sqrt{\pi}} = 0.26$

Elliptic flow: Freezeout?




switch from hydro to (weakly collisional) kinetics

at scale factor $b_{\perp}^{fr} = 1, 5, 10, 20$




no freezeout seen in the data


Elliptic flow: Shear vs bulk viscosity



Dissipative hydro with both η, ζ




Elliptic flow: Shear vs bulk viscosity



Dissipative hydro with both η, ζ

$$\beta_{\eta,\zeta} = (\eta, \zeta) \frac{E_F}{E} \frac{1}{(3\lambda N)^{1/3}}$$





$\eta \gg \zeta$

Dusling, Schaefer (2010)

Viscosity to entropy density ratio

consider both collective modes (low T)
and elliptic flow (high T)



Cao et al., Science (2010)

$$\eta/s \leq 0.4$$

Outlook

Experimental determination of transport properties: Collective modes and elliptic flow give $\langle \eta/s \rangle \lesssim 0.4$.

Local analysis requires second order hydro or hydro+kinetic.
(I am working on this.)

Shear viscous relaxation time can be measured by comparing collective modes and elliptic flow.

Can we observe breaking of scale invariance and the return of bulk viscosity away from unitarity? Can we measure η and ζ_3 in the superfluid phase?

Note: Experiment (Helium vs Fermi gas at unitarity)

