


# Bulk viscosity, spectra, and flow in heavy ion collisions

Thomas Schaefer & Kevin Dusling, North Carolina State University




# Why bulk viscosity?



Real QCD is not scale invariant, and  $\zeta \neq 0$ . Usually, this is treated as a nuisance – it leads to uncertainties in the extraction of  $\eta$ . Here, I want to estimate  $\zeta$  from data and see what (if anything) we can learn.


## Viscosity and dissipative forces

Shear viscosity determines shear stress (“friction”) in fluid flow



$$F = A \eta \frac{\partial v_x}{\partial y}$$


Bulk viscosity controls non-equilibrium pressure



$$P = P_0 - \zeta(\partial \cdot v)$$

# Shear and bulk viscosity in heavy ion collisions (first guess)

---



$$E_p \left. \frac{dN}{d^3p} \right|_{p_z=0} = v_0(p_\perp) (1 + 2v_2(p_\perp) \cos(2\phi) + \dots)$$

$\eta$  suppresses  $v_2$ , enhances  $v_0$

$\zeta$  suppresses  $v_0$ , (typically) enhances  $v_2$

Note:  $v_0$  also sensitive to eos, freezeout, hadronic phase.

# Differential elliptic flow from dissipative hydrodynamics



Spectra computed on freeze-out surface (“Cooper-Frye”)

$$E_p \frac{dN}{d^3p} = \frac{1}{(2\pi)^3} \int_{\sigma} f(E_p) p^{\mu} d\sigma_{\mu}$$

Write  $f = f^0 + \delta f$  and match to hydrodynamics

$$\delta\Pi^{\mu\nu} = \int d\Omega_p p^{\mu} p^{\nu} \delta f(E_p)$$

Only moments of  $\delta f$  fixed by  $\eta, \zeta$ . Need kinetic models.



## Relaxation time approximation

Approximate collision term by single relaxation time

$$C[\delta f_p] \simeq \frac{\delta f_p}{\tau(E_p)} \quad f_p = n_p^0 + \delta f_p$$

Bulk viscosity second order in conformal breaking parameter  $\delta c_s^2$

$$\zeta = 15\eta \left( c_s^2 - \frac{1}{3} \right)^2$$

Weinberg (1972)


Distribution function is first order in conformal breaking

$$\delta f \sim f_p^0 \frac{\eta}{sT} \frac{p^2}{T^2} \left( c_s^2 - \frac{1}{3} \right) (\partial \cdot u)$$


Near conformal fluids: Bulk viscous correction dominated by  $\delta f$

# Bulk viscosity in kinetic theory

QCD: Elastic vs inelastic reactions




$$g + g \rightarrow g + g \quad (m_g^2 \sim g^2 T^2)$$




$$g + g \rightarrow g + g + g$$

Hadron gas: inelastic scattering, hadro-chemistry



$$\pi + \pi \rightarrow 4\pi$$



$$p + \bar{p} \rightarrow 5\pi$$

## Distribution function in QGP

elastic  $2 \leftrightarrow 2$  can be written as Fokker-Planck equation (diffusion equation in momentum space)

$$(\partial \cdot u) \left( \frac{p^2}{3} - c_s^2 E_p \frac{\partial (\beta E_p)}{\partial \beta} \right) = \frac{T \mu_A}{n_p} \frac{\partial}{\partial p^i} \left( n_p \frac{\partial}{\partial p^i} \left[ \frac{\delta f_p}{n_p} \right] \right) + \dots$$


$$\text{drag coefficient } \mu_A = \frac{g^2 C_A m_D^2}{8\pi} \log \left( \frac{T}{m_D} \right)$$

Find  $\chi_B \sim \left( \frac{1}{3} - c_s^2 \right) \chi_S$  and (pure glue)


$$\zeta = \frac{0.44 \alpha_s^2 T^3}{\log(\alpha_s^{-1})} \quad \zeta \sim 47.9 \left( \frac{1}{3} - c_s^2 \right)^2 \eta$$



# Distribution function in QGP



Pure glue: shear vs bulk  
(bulk rescaled by  $\delta c_s^2$ )



QGP: quarks vs gluons

$$\delta f_p = -n_p(1 \pm n_p) [\chi_S(p) \hat{p}_i \hat{p}_j \sigma_{ij} + \chi_B(p) (\partial \cdot u)]$$

## Pion gas

Pion gas: Bulk viscosity governed by chemical non-equilibration

$$\delta f_p = n_p(1 + n_p) \left( \frac{\delta\mu}{T} + \frac{E_p \delta T}{T^2} \right) = -n_p(1 + n_p)(\chi_0 + \chi_1 E_p)(\partial \cdot u)$$

More formal:  $\chi_0$  is a “quasi zero mode” which dominates  $C^{-1}$

Inelastic rate determines  $\chi_0$ , energy conservation fixes  $\chi_1$

$$\chi_0 = \frac{\zeta}{\mathcal{F}} \quad \zeta = \frac{\beta \mathcal{F}^2}{4\Gamma_{2\pi \rightarrow 4\pi}}$$

where we have defined  $\mathcal{F} = \int d\Omega_p \left( \frac{p^2}{3} - c_s^2 E_p \frac{\partial(\beta E_p)}{\partial\beta} \right) n_p(1 + n_p)$

$$\zeta \simeq 12285 \frac{f_\pi^8}{m_\pi^5} \exp\left(-\frac{2m_\pi}{T}\right)$$

## Hadron resonance gas (model)

Hadron gas: Assume bulk viscosity dominated by chemical relaxation

$$\delta f_p^a = -n_p(1 \pm n_p) (\chi_0^a - \chi_1 E_p) (\partial \cdot u)$$

$\chi_0^a$  determined by rates,  $\chi_1$  fixed by energy conservation


Slowest rate determines  $\zeta$ , other rates fix  $\delta\mu^a/\delta\mu_\pi$ . Simple model

$$\chi_0^a \simeq \chi_0^\pi \begin{cases} 2 & \text{mesons} \\ 2.5 & \text{baryons} \end{cases}$$


inspired by  $\mu_\rho = 2\mu_\pi$  and  $2\mu_N = 5\mu_\pi$ . Find

$$\zeta/s = 0.05 \quad \Leftrightarrow \quad \delta\mu_\pi = 20 \text{ MeV}$$

# Bounds on $\zeta/s$ from differential $v_2$ (here: $K_s$ )




# Pion/Proton $p_T$ spectra (low $P_T$ )



Data: PHENIX nucl-ex/0307022. Hydro fit: Kevin Dusling (2012)

# Pion/Proton differential $v_2(p_T)$ spectra (low $p_T$ )



Data: STAR, nucl-ex/0409033. Hydro fit: Kevin Dusling (2012)

## Conclusions

Bulk viscous corrections dominated by freezeout distributions

QGP:  $\zeta$  controlled by momentum rearrangement


Hadron gas:  $\zeta$  determined by chemical non-equilibration

A new way to look at fugacity factors in thermal fits?


RHIC spectra seem to require  $\zeta/s \lesssim 0.05$

Bulk viscosity not zero: Spectra prefer  $\delta\mu$ , fine structure of  $v_2$  improves

## Extras: Second order hydrodynamics




gradient expansion  
(bulk stress)




freeze out surface  
(w/o bulk viscosity)



# Spectra and flow: Kaons and Lambdas




$$\eta/s = 0.16$$





$$\zeta/s = 0.04$$

# Flow: Interplay between shear and bulk viscosity



# Integrated $v_2$ versus centrality



## Distribution functions: Signs

Consider four-velocity  $u_\alpha$  with  $u^2 = -1$  ( $g_{\alpha\beta} = (-1, 1, 1, 1)$ )

$$\delta f_p = -n_p \chi_S p^\alpha p^\beta \langle \partial_\alpha u_\beta \rangle - n_p \chi_B (\partial \cdot u)$$

Asymptotic behavior  $\chi_{S,B} \sim p^2$ .


Consider BJ flow:  $p^\alpha p^\beta \langle \partial_\alpha u_\beta \rangle \sim -\frac{p_T^2}{\tau}$  and  $\partial \cdot u \sim \frac{1}{\tau}$ .

$$\delta f_p \sim \frac{\eta}{s} \left( \frac{p_T}{T} \right)^2 \frac{1}{\tau T} - \frac{\zeta}{s} \left( \frac{p_T}{T} \right)^2 \frac{1}{\tau T}$$


Elliptic flow

$$\langle v_2 \rangle = \frac{\int d\phi [f(\phi) + \delta f(\phi)] \cos(2\phi)}{\int d\phi [f(\phi) + \delta f(\phi)]} \simeq \langle v_2^0 \rangle + \langle \delta v_2 \rangle - \langle v_2^0 \rangle \langle \delta v_0 \rangle$$

# Elliptic flow: Shear vs bulk viscosity





Dissipative hydro with both  $\eta, \zeta$



# Elliptic flow: Shear vs bulk viscosity

Dissipative hydro with both  $\eta, \zeta$

$$\beta_{\eta, \zeta} = (\eta, \zeta) \frac{E_F}{E} \frac{1}{(3\lambda N)^{1/3}}$$



$\eta \gg \zeta$