

Nearly Perfect Fluidity


in Cold Atomic Gases

Thomas Schaefer, North Carolina State University




Fluids: Gases, liquids, plasmas, . . .

Hydrodynamics: Long-wavelength, low-frequency dynamics of conserved or spontaneously broken symmetry variables.




$$\tau \sim \tau_{micro}$$



$$\tau \sim \lambda$$

Historically: Water
 $(\rho, \epsilon, \vec{\pi})$



Simple non-relativistic fluid

Simple fluid: Conservation laws for mass, energy, momentum

$$\frac{\partial \rho}{\partial t} + \vec{\nabla}(\rho \vec{v}) = 0$$

$$\frac{\partial \epsilon}{\partial t} + \vec{\nabla} \vec{j}^\epsilon = 0$$

$$\frac{\partial}{\partial t}(\rho v_i) + \frac{\partial}{\partial x_j} \Pi_{ij} = 0$$



Constitutive relations: Energy momentum tensor

$$\Pi_{ij} = P\delta_{ij} + \rho v_i v_j + \eta \left(\partial_i v_j + \partial_j v_i - \frac{2}{3} \delta_{ij} \partial_k v_k \right) + O(\partial^2)$$

reactive

dissipative

2nd order

$$\text{Expansion } \Pi_{ij}^0 \gg \delta \Pi_{ij}^1 \gg \delta \Pi_{ij}^2$$

Regime of applicability

Expansion parameter $Re^{-1} = \frac{\eta(\partial v)}{\rho v^2} = \frac{\eta}{\rho Lv} \ll 1$


$$Re^{-1} = \frac{\eta}{\hbar n} \times \frac{\hbar}{mvL}$$

fluid flow
property property

Consider $mvL \sim \hbar$: Hydrodynamics requires $\eta/(\hbar n) < 1$

Shear viscosity

Viscosity determines shear stress (“friction”) in fluid flow




$$F = A \eta \frac{\partial v_x}{\partial y}$$

Kinetic theory: conserved quantities carried by quasi-particles

$$\frac{\partial f_p}{\partial t} + \vec{v} \cdot \vec{\nabla}_x f_p + \vec{F} \cdot \vec{\nabla}_p f_p = C[f_p]$$

$$\eta \sim \frac{1}{3} n \bar{p} l_{mfp}$$



Dilute, weakly interacting gas: $l_{mfp} \sim 1/(n\sigma)$

$$\eta \sim \frac{1}{3} \frac{\bar{p}}{\sigma}$$

independent of density!

Shear viscosity

non-interacting gas ($\sigma \rightarrow 0$):

$$\eta \rightarrow \infty$$


non-interacting and hydro limit ($T \rightarrow \infty$) limit do not commute

strongly interacting gas:

$$\frac{\eta}{n} \sim \bar{p} l_{mfp} \geq \hbar$$

but: kinetic theory not reliable!

what happens if the gas condenses into a liquid?



Eyring, Frenkel:

$$\eta \simeq hn \exp(E/T) \geq hn$$


Holographic duals: Transport properties

Thermal (conformal) field theory $\equiv AdS_5$ black hole

| | | |
|-----------------|-------------------|---|
| CFT temperature | \Leftrightarrow | Hawking temperature |
| CFT entropy | \Leftrightarrow | Hawking-Bekenstein entropy \sim area of event horizon |
| shear viscosity | \Leftrightarrow | Graviton absorption cross section \sim area of event horizon |

$$T_{\mu\nu} = \frac{1}{\sqrt{-g}} \frac{\delta S}{\delta g_{\mu\nu}}$$

$$g_{\mu\nu} = g_{\mu\nu}^0 + \gamma_{\mu\nu}$$



Holographic duals: Transport properties

Thermal (conformal) field theory $\equiv AdS_5$ black hole

CFT entropy \Leftrightarrow

Hawking-Bekenstein entropy

\sim area of event horizon

shear viscosity \Leftrightarrow


Graviton absorption cross section

\sim area of event horizon

Strong coupling limit

$$\frac{\eta}{s} = \frac{\hbar}{4\pi k_B}$$


Son and Starinets (2001)




Strong coupling limit universal? Provides lower bound for all theories?

Answer appears to be no; e.g. theories with higher derivative gravity duals.

Kinetics vs No-Kinetics



AdS/CFT low viscosity goo



pQCD kinetic plasma

Effective theories for fluids (Here: Weak coupling QCD)



$$\mathcal{L} = \bar{q}_f(iD - m_f)q_f - \frac{1}{4}G_{\mu\nu}^a G_{\mu\nu}^a$$



$$\frac{\partial f_p}{\partial t} + \vec{v} \cdot \vec{\nabla}_x f_p = C[f_p] \quad (\omega < T)$$



$$\frac{\partial}{\partial t}(\rho v_i) + \frac{\partial}{\partial x_j}\Pi_{ij} = 0 \quad (\omega < g^4 T)$$

Effective theories (Strong coupling)



$$\mathcal{L} = \bar{\lambda}(i\sigma \cdot D)\lambda - \frac{1}{4}G_{\mu\nu}^a G_{\mu\nu}^a + \dots \Leftrightarrow S = \frac{1}{2\kappa_5^2} \int d^5x \sqrt{-g} \mathcal{R} + \dots$$



$$\frac{\partial}{\partial t}(\rho v_i) + \frac{\partial}{\partial x_j} \Pi_{ij} = 0 \quad (\omega < T)$$

Perfect Fluids: How to be a contender?

Bound is quantum mechanical

need quantum fluids



Bound is incompatible with weak coupling and kinetic theory

strong interactions, no quasi-particles

Model system has conformal invariance (essential?)

(Almost) scale invariant systems

Perfect Fluids: The contenders




QGP ($T=180$ MeV)

trapped atoms
($T=0.1$ neV)




Liquid Helium
($T=0.1$ meV)

Perfect Fluids: The contenders



QGP $\eta = 5 \cdot 10^{11} Pa \cdot s$



Trapped Atoms

$\eta = 1.7 \cdot 10^{-15} Pa \cdot s$



Liquid Helium


$\eta = 1.7 \cdot 10^{-6} Pa \cdot s$

Consider ratios

η/s


Unitarity limit

Consider simple square well potential




$$a < 0$$

$$a = \infty, \epsilon_B = 0$$




$$a > 0, \epsilon_B > 0$$



Unitarity limit

Now take the range to zero, keeping $\epsilon_B \simeq 0$



Universal relations

$$\mathcal{T} = \frac{1}{ik + 1/a}$$

$$\epsilon_B = \frac{1}{2ma^2}$$

$$\psi_B \sim \frac{1}{\sqrt{ar}} \exp(-r/a)$$

Dilute Fermi gas: field theory

Non-relativistic fermions at low momentum

$$\mathcal{L}_{\text{eff}} = \psi^\dagger \left(i\partial_0 + \frac{\nabla^2}{2M} \right) \psi - \frac{C_0}{2} (\psi^\dagger \psi)^2$$

Unitary limit: $a \rightarrow \infty, \sigma \rightarrow 4\pi/k^2$ ($C_0 \rightarrow \infty$)

This limit is smooth: HS-trafo, $\Psi = (\psi_\uparrow, \psi_\downarrow^\dagger)$

$$\mathcal{L} = \Psi^\dagger \left[i\partial_0 + \sigma_3 \frac{\vec{\nabla}^2}{2m} \right] \Psi + (\Psi^\dagger \sigma_+ \Psi \phi + h.c.) - \frac{1}{C_0} \phi^* \phi ,$$

Low T ($T < T_c \sim \mu$): Pairing and superfluidity, $\langle \phi \rangle \neq 0$

Linear response and kinetic theory

Consider background metric $g_{ij}(t, \mathbf{x}) = \delta_{ij} + h_{ij}(t, \mathbf{x})$. Linear response


$$\delta\Pi^{ij} = \frac{\delta\Pi_{ij}^{eq}}{\delta h_{ij}} h^{ij} - \frac{1}{2} G_R^{ijkl} h_{kl}$$

Kubo relation: $\eta(\omega) = \frac{1}{\omega} \text{Im}G_R^{xyxy}(\omega, 0)$

Kinetic theory: Boltzmann equation

$$\left(\frac{\partial}{\partial t} + \frac{p^i}{m} \frac{\partial}{\partial x^i} - \left(g^{il} \dot{g}_{lj} p^j + \Gamma_{jk}^i \frac{p^j p^k}{m} \right) \frac{\partial}{\partial p^i} \right) f(t, \mathbf{x}, \mathbf{p}) = C[f]$$

$$C[f] =$$




Kinetic theory

linearize $f = f_0 + \delta f$, solve for δf , $\hookrightarrow \delta \Pi_{ij}$, $\hookrightarrow G_R$, $\hookrightarrow \eta(\omega)$


$$\eta(\omega) = \frac{\eta}{1 + \omega^2 \tau_R^2}$$

$$\eta = \frac{15}{32\sqrt{\pi}} (mT)^{3/2} \quad \tau_R = \frac{\eta}{nT}$$

shear channel



sound channel



Shear viscosity: Sum rules



Randeira and Taylor proved the following sum rules

$$\frac{1}{\pi} \int dw \left[\eta(\omega) - \frac{C}{10\pi\sqrt{m\omega}} \right] = \frac{\varepsilon}{3} - \frac{C}{10\pi ma}$$



$$\frac{1}{\pi} \int dw \zeta(\omega) = \frac{1}{72\pi ma^2} \left(\frac{\partial C}{\partial a^{-1}} \right)$$

where C is Tan's contact, $\rho(k) \sim C/k^4$.


Sum rules constrain spectral function and euclidean correlator



Almost ideal fluid dynamics




Hydrodynamic
expansion converts
coordinate space
anisotropy
to momentum space
anisotropy



Hydrodynamics: Collective modes

Radial breathing mode



Ideal fluid hydrodynamics ($P = \frac{2}{3}\mathcal{E}$)

$$\begin{aligned} \frac{\partial n}{\partial t} + \vec{\nabla} \cdot (n\vec{v}) &= 0 \\ \frac{\partial \vec{v}}{\partial t} + (\vec{v} \cdot \vec{\nabla}) \vec{v} &= -\frac{\vec{\nabla} P}{mn} - \frac{\vec{\nabla} V}{m} \end{aligned}$$

Hydro frequency at unitarity

$$\omega = \sqrt{\frac{10}{3}} \omega_\perp$$

Damping small, depends on T/T_F .


Damping of collective mode

Energy dissipation (η, ζ, κ : shear, bulk viscosity, heat conductivity)

$$\begin{aligned}\dot{E} = & -\frac{1}{2} \int d^3x \eta(x) \left(\partial_i v_j + \partial_j v_i - \frac{2}{3} \delta_{ij} \partial_k v_k \right)^2 \\ & - \int d^3x \zeta(x) (\partial_i v_i)^2 - \frac{1}{T} \int d^3x \kappa(x) (\partial_i T)^2\end{aligned}$$

Shear viscosity to entropy ratio
(assuming $\zeta = \kappa = 0$)





$$\frac{\eta}{s} = (3\lambda N)^{\frac{1}{3}} \frac{\Gamma}{\omega_{\perp}} \frac{E_0}{E_F} \frac{N}{S}$$



Schaefer (2007), see also Bruun, Smith

$T \ll T_F$ $T \gg T_F, \tau_R \simeq \eta/P$

Hydrodynamics: Free expansion and rotation



Scaling Flows

Universal equation of state

$$P = \frac{2}{3}\mathcal{E}$$

Equilibrium density profile

$$n_0(x) = n(\mu(x), T) \quad \mu(x) = \mu_0 \left(1 - \frac{x^2}{R_x^2} - \frac{y^2}{R_y^2} - \frac{z^2}{R_z^2} \right)$$

Scaling Flow: Stretch and rotate profile

$$\mu_0 \rightarrow \mu_0(t), \quad T \rightarrow T_0(\mu_0(t)/\mu_0), \quad R_x \rightarrow R_x(t), \dots$$

Linear velocity profile

$$\vec{v}(x, t) = (\alpha_x x, \alpha_y y, \alpha_z z) + \alpha \vec{\nabla}(xy) + \vec{\omega} \times \vec{x}$$

“Hubble flow”

Scaling hydrodynamics

Write $R_i(t) = b_i(t)R_i(0)$. Euler equation

$$\frac{\ddot{b}_\perp}{b_\perp} = \frac{\omega_\perp^2}{(b_\perp^2 b_{||})^{2/3}} \frac{1}{b_\perp^2} \quad b_\perp(\omega_\perp t \gg 1) \sim \sqrt{\frac{3}{2}} \omega_\perp t$$

Dissipation breaks scaling behavior ($\nabla_i P/n = a_i x_i$)



$$\frac{\ddot{b}_\perp}{b_\perp} = a_\perp - \frac{2\beta\omega_\perp}{b_\perp^2} \left(\frac{\dot{b}_\perp}{b_\perp} - \frac{\dot{b}_x}{b_x} \right) \quad friction$$

$$\dot{a}_\perp = ideal + \frac{8\beta\omega_\perp^2}{3b_\perp} \left(\frac{\dot{b}_\perp}{b_\perp} - \frac{\dot{b}_z}{b_z} \right)^2 \quad heating$$

$$\beta = \frac{\langle \eta \rangle}{N} \frac{E_F}{E_0} \frac{1}{(3N\lambda)^{1/3}}$$

Navier-Stokes: Numerical results

Full 3-D hydro with $\eta = \alpha_n n$ and $\alpha_n = const.$



Reheating leads to reacceleration. Dissipation causes characteristic curvature of $A_R(t) \equiv R_{\perp}/R_z$.

- Issues:
- i) Dilute corona $\eta \sim T^{3/2} \rightarrow \nabla_i \delta \Pi_{ij} = 0$. No force (?)
 - ii) $Kn \sim (b_{||}/b_{\perp})^{1/3}$ drops \rightarrow No freezeout (?)


Relaxation time model

In real systems stress tensor does not relax to Navier-Stokes form instantaneously. Consider


$$\tau_R \left(\frac{\partial}{\partial t} + \vec{v} \cdot \vec{\nabla} \right) \delta\Pi_{ij} = \delta\Pi_{ij} - \eta \left(\partial_i v_j + \partial_j v_i - \frac{2}{3} \delta_{ij} \partial \cdot v \right)$$

In kinetic theory $\tau_R \simeq (\eta/n) T^{-1}$


- dissipation from $\eta \sim (mT)^{3/2}$: corona exerts drag force.
- find $\langle \alpha_n \rangle \sim T^3$
- system dependence



Elliptic flow: High T limit




$$\text{Quantum viscosity } \eta = \eta_0 \frac{(mT)^{3/2}}{\hbar^2}$$



$$\text{fit: } \eta_0 = 0.33 \pm 0.04$$


$$\text{theory: } \eta_0 = \frac{15}{32\sqrt{\pi}} = 0.26$$

Elliptic flow: Freezeout?



switch from hydro to (weakly collisional) kinetics

at scale factor $b_{\perp}^{fr} = 1, 5, 10, 20$





no freezeout seen in the data

Elliptic flow: Shear vs bulk viscosity



Dissipative hydro with both η, ζ

$$\beta_{\eta,\zeta} = (\eta, \zeta) \frac{E_F}{E} \frac{1}{(3\lambda N)^{1/3}}$$



Viscosity to entropy density ratio

consider both collective modes (low T)
and elliptic flow (high T)



Cao et al., Science (2010)

$$\eta/s \leq 0.4$$

Outlook

The unitary Fermi gas is an important model system for other strongly correlated quantum fluids in nature.




The equation of state has been determined to a few percent.

Transport properties are more difficult: Kinetic theory at $T \gg T_F$ and $T \ll T_F$. Sum rules constrain spectral fct at all T .

Experimental determination of transport properties: Collective modes give $\langle \eta/s \rangle < 0.4$. Local analysis requires second order hydro or hydro+kinetic.

Extra

Kinetic theory summary



Experiment (Liquid Helium)





FIG. 1. The viscosity of liquid helium II measured by flow through a 10^{-4} cm channel.

Kapitza (1938)
viscosity vanishes below T_c
capillary flow viscometer





Hollis-Hallett (1955)
roton minimum, phonon rise
rotation viscometer

$$\eta/s \simeq 0.8 \hbar/k_B$$

Elliptic Flow (QGP)


Hydrodynamic expansion converts coordinate space anisotropy to momentum space anisotropy



source: U. Heinz (2005)

$$p_0 \left. \frac{dN}{d^3p} \right|_{p_z=0} = v_0(p_\perp) (1 + 2v_2(p_\perp) \cos(2\phi) + \dots)$$

Elliptic flow: initial entropy scaling



source: U. Heinz (2005)


Viscosity and Elliptic Flow

Consistency condition $T_{\mu\nu} \gg \delta T_{\mu\nu}$
(applicability of Navier-Stokes)

$$\frac{\eta + \frac{4}{3}\zeta}{s} \ll \frac{3}{4}(\tau T)$$

Danielewicz, Gyulassy (1985)

Very restrictive for $\tau < 1$ fm



Romatschke (2007), Teaney (2003)

Many questions: Dependence on initial conditions, freeze out, etc.

conservative bound

$$\frac{\eta}{s} < 0.4$$