

Effective Theories of Very Dense Matter: From keV to GeV

Thomas Schaefer

North Carolina State

w. K. Schwenzer (NCSU) and A. Kryjevski (INT)

Motivation

There is a successful effective theory of fermionic many body systems

Landau Fermi-Liquid Theory


FLT theory: Quasi-particles near the Fermi surface. Interactions characterized by Fermi-liquid parameters.

Predicts collective modes, thermodynamics, transport, . . .

Gauge Theories: Unscreened long range forces

Does a quasi-particle EFT exist?


Effective Field Theories



High Density Effective Theory

Quasi-particles (holes)


$$E_{\pm} = -\mu \pm \sqrt{\vec{p}^2 + m^2} \simeq -\mu \pm |\vec{p}|$$



Effective field theory on v -patches

$$\psi_{v\pm} = e^{-i\mu v \cdot x} \left(\frac{1 \pm \vec{\alpha} \cdot \vec{v}}{2} \right) \psi$$


Effective lagrangian for ψ_{v+}




$$\mathcal{L} = \sum_v \psi_v^\dagger (iv \cdot D) \psi_v - \frac{1}{4} G_{\mu\nu}^a G_{\mu\nu}^a + O(1/\mu)$$

Four Quark Operators

quark-quark scattering



BCS




Landau

$$\mathcal{L}_{BCS} = \frac{1}{\mu^2} \sum V_l^{\Gamma\Gamma'} R_l^{\Gamma\Gamma'} (\vec{v} \cdot \vec{v}') \left(\psi_v \Gamma \psi_{-v} \right) \left(\psi_{v'}^\dagger \Gamma' \psi_{-v'}^\dagger \right),$$

$$\mathcal{L}_{FL} = \frac{1}{\mu^2} \sum F_l^{\Gamma\Gamma'}(\phi) R_l^{\Gamma\Gamma'}(\vec{v} \cdot \vec{v}') \left(\psi_v \Gamma \psi_{v'} \right) \left(\psi_{\tilde{v}}^\dagger \Gamma' \psi_{\tilde{v}'}^\dagger \right)$$

Four Fermion Operators: Matching

Match scattering amplitudes on Fermi surface: forward scattering



Color-flavor-spin symmetric terms


$$f_0^s = \frac{C_F}{4N_c N_f} \frac{g^2}{p_F^2}, \quad f_i^s = 0 \quad (i > 1)$$

Power Counting

Naive power counting

$$\mathcal{L} = \hat{\mathcal{L}} \left(\psi, \psi^\dagger, \frac{D_{||}}{\mu}, \frac{D_\perp}{\mu}, \frac{\bar{D}_{||}}{\mu}, \frac{m}{\mu} \right)$$


Problem: hard loops



$$\frac{1}{2\pi} \sum_{\vec{v}} \int \frac{d^2 l_\perp}{(2\pi)^2} = \frac{\mu^2}{2\pi^2} \int \frac{d\Omega}{4\pi}.$$

Modified Power Counting: $\mathcal{A} \sim l^\delta$

$$\delta = \sum_i [(k-4)V_k^S + (k-2-f_k)V_k^H] + E_Q + 4 - 2N_C$$




V_k^S soft vertices of $O(l^k)$

V_k^H hard vertices of $O(l^k)$

f_k number of fermion fields in V_k^H

E_Q external quark lines

N_C connected graphs in hard graph



- resum quark loops in gluon n-pt fcts
- four quark operators are leading order
- six quark operators (etc) are suppressed

Effective Theory for $l \sim g\mu$

$$\mathcal{L} = \psi_v^\dagger (iv \cdot D) \psi_v + f_0^s (\psi_v^\dagger \psi_v) (\psi_{v'}^\dagger \psi_{v'}) - \frac{1}{4} G_{\mu\nu}^a G_{\mu\nu}^a + \mathcal{L}_{HDL}$$

$$\mathcal{L}_{HDL} = -\frac{m^2}{2} \sum_v G_{\mu\alpha}^a \frac{v^\alpha v^\beta}{(v \cdot D)^2} G_{\mu\beta}^b$$


Transverse gauge boson propagator

$$D_{ij}(k) = \frac{\delta_{ij} - \hat{k}_i \hat{k}_j}{k_0^2 - \vec{k}^2 + i\eta|k_0|/|\vec{k}|},$$

Scaling of gluon momenta

$$|\vec{k}| \sim k_0^{1/3} \eta^{2/3} \gg k_0 \quad \text{gluons are very spacelike}$$


Effective Theory for $l < g\mu$




Low energy expansion contains fractional powers $\omega^{1/3}$

Quasi-Quarks at Large Density: Non-Fermi Liquid Effects

Loop Corrections



$$\Sigma(\omega) \simeq \frac{g^2 C_F}{12\pi^2} \omega \log \left(\frac{\Lambda}{\omega} \right)$$



$$\Gamma_\alpha = \frac{g^3 C_F v_\alpha}{12\pi^2} \log \left(\frac{\Lambda}{\omega} \right)$$

$$\Gamma_\alpha = O(g^3)$$

time
like

space
like

“Migdal’s Theorem” for QCD

- self energy non-perturbative (?) for $\omega \sim \Lambda \exp(-9\pi^2/g^2)$
- no large logarithms in vertex ($k \gg k_4!$)

Renormalization Group

Renormalized parameters

$$\psi_{0,v} = Z^{1/2} \psi_v, \quad v_{0,F} = Z_F v_F, \quad g_0 = \frac{Z_g}{ZZ_F} g, \quad \alpha = \frac{g^2 v_F}{4\pi},$$

One-loop calculation ($\beta = \frac{\partial \alpha}{\partial \log \Lambda}$, $\gamma = \frac{\partial \log Z}{\partial \log \Lambda}$, $\gamma_F = \frac{\partial \log Z_F}{\partial \log \Lambda}$)

$$\gamma(\alpha) = -\gamma_F(\alpha) = \frac{4\alpha}{9\pi}, \quad \beta(\alpha) = -\gamma_F(\alpha)\alpha \quad \text{IR free !!}$$

RG equation

$$\left\{ \Lambda \frac{\partial}{\partial \Lambda} + \beta(\alpha) \frac{\partial}{\partial \alpha} - \gamma_F(\alpha) l_i \frac{\partial}{\partial l_i} + \frac{n}{2} \gamma(\alpha) \right\} G^{(n)}(\omega_i, l_i, \alpha) = 0,$$

$\gamma \ll 1$: RG equation can be solved exactly

$$S^{-1}(\omega, l) = \omega \left(1 + \gamma \log \left(\frac{\Lambda}{\omega} \right) \right) - v_F l$$

no terms $\alpha^2 \log^2(\omega)$, etc.

- quasi-particle velocity vanishes as


$$v \sim \log(\Lambda/\omega)^{-1}$$

- anomalous term in the specific heat

$$c_v \sim \gamma T \log(T)$$

- enhanced corrections to the gap

$$\log(\mu/\Delta) = \log(\mu/\Delta_0)(1 - O(\gamma g))$$



Remnants of Fermi Liquid Behavior

Perturbative corrections to matching QCD→HDET

$$v_F = 1 - \frac{C_F \alpha_s}{2\pi}, \quad \delta\mu = \frac{C_F \alpha_s}{\pi} \mu.$$

These coefficients satisfy Fermi-liquid relations (Baym&Chin)



$$v_F = \frac{p_F}{\mu} - \frac{N f_1^s}{3} \quad p_F = \mu \left(1 - \frac{N f_0^s}{2} \right)$$

Fermi-liquid parameters



$$f_0^s = \frac{C_F}{4N_c N_f} \frac{g^2}{p_F^2} \quad f_1^s = 0$$

Application: Neutrino Emission

Quark Direct URCA: $d \rightarrow u + e^- + \bar{\nu}$, $u + e^- \rightarrow d + \bar{\nu}$



$$\epsilon \sim G_F T^7$$




$$\epsilon \sim G_F \alpha_s^3 T^6 \log^2(T)$$

Quasi-Baryons at Large Density: CFL Phase

Superconductivity

quark-quark scattering

$$(\mu \gg \Lambda_{QCD})$$



Gap equation: double logarithmic behavior

$$\Delta(p_0) = \frac{g^2}{18\pi^2} \int dq_0 \left\{ \log \left(\frac{b_M}{|p_0 - q_0|} \right) + \dots \right\} \frac{\Delta(q_0)}{\sqrt{q_0^2 + \Delta(q_0)^2}}$$

collinear log

BCS log

$$\Rightarrow \quad \Delta_0 = 512\pi^4 \mu g^{-5} \exp\left(-\frac{\pi^2 + 4}{8}\right) \exp\left(-\frac{3\pi^2}{\sqrt{2}g}\right)$$

CFL Phase

Consider $N_f = 3$ ($m_i = 0$)

$$\langle q_i^a q_j^b \rangle = \phi \epsilon^{abI} \epsilon_{ijI}$$


$$\langle ud \rangle = \langle us \rangle = \langle ds \rangle$$

$$\langle rb \rangle = \langle rg \rangle = \langle bg \rangle$$

Symmetry breaking pattern:

$$SU(3)_L \times SU(3)_R \times [SU(3)]_C \\ \times U(1) \rightarrow SU(3)_{C+F}$$

All quarks and gluons acquire a gap



$$\langle \psi_L \psi_L \rangle = -\langle \psi_R \psi_R \rangle$$

EFT in the CFL Phase

Consider HDET with a CFL gap term

$$\mathcal{L} = \text{Tr} \left(\psi_L^\dagger (iv \cdot D) \psi_L \right) + \frac{\Delta}{2} \left\{ \text{Tr} \left(X^\dagger \psi_L X^\dagger \psi_L \right) - \kappa \left[\text{Tr} \left(X^\dagger \psi_L \right) \right]^2 \right\}$$

$$+ (L \leftrightarrow R, X \leftrightarrow Y)$$

$$\psi_L \rightarrow L \psi_L C^T, \quad X \rightarrow L X C^T, \quad \langle X \rangle = \langle Y \rangle = \mathbb{1}$$

Quark loops generate a kinetic term for X, Y

$$\mathcal{L} = -\frac{f_\pi^2}{2} \left\{ \text{Tr} \left((X^\dagger D_0 X)^2 + (Y^\dagger D_0 Y)^2 \right) \right\} + \dots$$

Integrate out gluons, identify low energy fields ($\xi = \Sigma^{1/2}$)

$$\Sigma = XY^\dagger \quad [8] + [1] \text{ GBs}, \quad N_L = \xi (\psi_L X^\dagger) \xi^\dagger \quad [8] + [1] \text{ Baryons}$$

Effective theory: CFL(B)χPTh

$$\begin{aligned}\mathcal{L} = & \frac{f_\pi^2}{4} \left\{ \text{Tr} (\nabla_0 \Sigma \nabla_0 \Sigma^\dagger) - v_\pi^2 \text{Tr} (\nabla_i \Sigma \nabla_i \Sigma^\dagger) \right\} \\ & + \text{Tr} (N^\dagger i v^\mu D_\mu N) - D \text{Tr} (N^\dagger v^\mu \gamma_5 \{\mathcal{A}_\mu, N\}) \\ & - F \text{Tr} (N^\dagger v^\mu \gamma_5 [\mathcal{A}_\mu, N]) + \frac{\Delta}{2} \left\{ \text{Tr} (NN) - [\text{Tr} (N)]^2 \right\}\end{aligned}$$

with $D_\mu N = \partial_\mu N + i[\mathcal{V}_\mu, N]$


$$\begin{aligned}\mathcal{V}_\mu &= -\frac{i}{2} (\xi \partial_\mu \xi^\dagger + \xi^\dagger \partial_\mu \xi) \\ \mathcal{A}_\mu &= -\frac{i}{2} \xi (\partial_\mu \Sigma^\dagger) \xi\end{aligned}$$

$$f_\pi^2 = \frac{21 - 8 \log 2}{18} \frac{\mu^2}{2\pi^2} \quad v_\pi^2 = \frac{1}{3} \quad D = F = \frac{1}{2}$$

Mass Terms: Match HDET to QCD

$$\mathcal{L} = \psi_R^\dagger \frac{MM^\dagger}{2\mu} \psi_R + \psi_L^\dagger \frac{M^\dagger M}{2\mu} \psi_L$$

$$+ \frac{C}{\mu^2} (\psi_R^\dagger M \lambda^a \psi_L) (\psi_R^\dagger M \lambda^a \psi_L)$$



mass corrections to FL parameters $\hat{\mu}$ and $F^0(++) \rightarrow --$)

Mass Terms: Match HDET to CFL χ Th

Kinetic term: $\psi_L^\dagger X_L \psi_L + \psi_R^\dagger X_R \psi_R$

$$\begin{aligned} D_0 N &= \partial_0 N + i[\Gamma_0, N], \quad \Gamma_0 = \mathcal{V}_0 + \frac{1}{2} (\xi X_R \xi^\dagger + \xi^\dagger X_L \xi) \\ \nabla_0 \Sigma &= \partial_0 \Sigma + iX_L \Sigma - i\Sigma X_R \end{aligned}$$

vector (axial) potentials

Contact term: $(\psi_R^\dagger M \psi_L)(\psi_R^\dagger M \psi_L)$

$$\mathcal{L} = \frac{3\Delta^2}{4\pi^2} \left\{ [\text{Tr}(M\Sigma)]^2 - \text{Tr}(M\Sigma M\Sigma) \right\}$$

meson mass terms

Phase Structure and Spectrum

Phase structure determined by effective potential

$$V(\Sigma) = \frac{f_\pi^2}{2} \text{Tr} (X_L \Sigma X_R \Sigma^\dagger) - A \text{Tr}(M \Sigma^\dagger) - B_1 [\text{Tr}(M \Sigma^\dagger)]^2 + \dots$$


$$V(\Sigma_0) \equiv \min$$

Fermion spectrum determined by

$$\mathcal{L} = \text{Tr} (N^\dagger i v^\mu D_\mu N) + \text{Tr} (N^\dagger \gamma_5 \rho_A N) + \frac{\Delta}{2} \left\{ \text{Tr} (NN) - [\text{Tr} (N)]^2 \right\},$$


$$\rho_{V,A} = \frac{1}{2} \left\{ \xi \frac{M^\dagger M}{2p_F} \xi^\dagger \pm \xi^\dagger \frac{M M^\dagger}{2p_F} \xi \right\} \quad \xi = \sqrt{\Sigma_0}$$

Phase Structure and Spectrum



meson condensation: CFLK

reliable: yes!



gapless modes? (gCFLK)

reliable: not clear yet

Summary

EFT/RG methods provide powerful tools

phase structure and spectrum at large density

Normal phase: Non-Fermi liquid behavior due to unscreened transverse gauge bosons

modified quasi-particles

but: no break-down of perturbation theory

Superfluid phase: effective chiral theory with calculable coefficients

kaon condensation, possibility of gapless modes