

## Ideal Gas

Ideal gas law

$$PV = kNT$$

Equipartition law

$$U = \frac{f}{2}kNT$$

with  $f = 3$  for a mono-atomic gas and  $f = 5$  for a di-atomic gas. Adiabatic expansion

$$pV^\gamma = \text{const}, \quad \gamma = (f + 2)/f$$

Entropy of an ideal mono-atomic gas

$$S = kN \left\{ \log \left( \frac{V}{Nv_Q} \right) + \frac{5}{2} \right\}, \quad v_Q = l_Q^3, \quad l_Q = \frac{h}{\sqrt{2\pi mkT}}$$

Chemical potential

$$\mu = -kT \log \left( \frac{V}{Nv_Q} \right)$$

## Entropy and Heat

First law

$$\Delta U = Q + W$$

Thermodynamic Identity

$$dU = TdS - PdV + \mu dN$$

If  $W = -PdV$  have  $Q = TdS$ . Also

$$\frac{1}{T} = \left. \frac{\partial S}{\partial U} \right|_{V,N}, \quad P = T \left. \frac{\partial S}{\partial V} \right|_{U,N}$$

Specific heat  $C = Q/\Delta T$ . Have

$$C_V = \left. \frac{\partial U}{\partial T} \right|_{V,N}$$

Efficiency of the Carnot Process operating between two reservoirs at temperatures  $T_h$  and  $T_c$

$$\epsilon = \frac{W}{Q_h} = 1 - \frac{T_c}{T_h}$$

## Thermodynamic Functions

Enthalpy

$$H = U + PV \quad \Delta H = Q + W_{other} \quad (P = const)$$

Free Energy

$$F = U - TS \quad \Delta F = W \quad (T = const, Q = T\Delta S)$$

Gibbs Free Energy

$$G = U - TS + PV \quad \Delta G = W_{other} \quad (P = T = const, Q = T\Delta S)$$

Partial derivatives of free energy

$$S = - \left( \frac{\partial F}{\partial T} \right)_{V,N} \quad P = - \left( \frac{\partial F}{\partial V} \right)_{T,N} \quad \mu = \left( \frac{\partial F}{\partial N} \right)_{V,T}$$

## Statistical Definition of Entropy

Entropy

$$S = k \log(\Omega)$$

Binomial coefficient

$$\binom{N}{k} = \frac{N!}{k!(N-k)!}$$

Stirling formula ( $N \gg 1$ )

$$\log(N!) \simeq N \log(N) - N + \dots$$

## Statistical Mechanics

Partition Function

$$Z = \sum_s \exp(-\beta E_s), \quad \beta = \frac{1}{kT}$$

Average (internal) energy

$$\bar{E} = -\frac{1}{Z} \frac{\partial Z}{\partial \beta}$$

Connection to free energy

$$F = -kT \log(Z)$$

For  $N$  not-to-dense, indistinguishable particles

$$Z_{tot} = \frac{1}{N!} (Z_1)^N,$$

where  $Z_1$  is the one-body partition function. Grand partition Function

$$\mathcal{Z} = \sum_s \exp(-\beta(E_s - \mu N_s))$$

Bose and Fermi distribution

$$n_B = \frac{1}{\exp(\beta(\epsilon - \mu)) - 1}, \quad n_F = \frac{1}{\exp(\beta(\epsilon - \mu)) + 1}$$

Boltzmann limit  $n = \exp(-\beta(E - \mu))$

## Numerical Constants

$$\begin{aligned} k &= 1.381 \times 10^{-23} \text{ J/K} = 8.617 \times 10^{-5} \text{ eV/K} \\ N_A &= 6.022 \times 10^{23} \\ R &= 8.315 \text{ J/mol/K} \\ h &= 6.626 \times 10^{-34} \text{ J} \cdot \text{s} \\ 1 \text{ atm} &= 1.013 \times 10^5 \text{ N/m}^2 \\ 1 \text{ cal} &= 4.186 \text{ J} \\ 1 \text{ eV} &= 1.602 \times 10^{-19} \text{ J} \\ 1 \text{ u} &= 1.661 \times 10^{-27} \text{ kg} \end{aligned}$$